Approximate Calculation of Transport Coefficients of Earth and Mars Atmospheric Dissociating Gases

Vladimir V. Riabov*

Worcester Polytechnic Institute, Worcester, Massachusetts 01609-2280

High-temperature transport properties (viscosity, thermal conductivity, binary and multicomponent mass diffusion, and thermal diffusion) of dissociating gases of the Earth and Martian atmospheres have been calculated within the framework of the Chapman-Enskog method. The exponential repulsion potentials of interactions between atoms (N, O, and C) and molecules (N₂, O₂, NO, C₂, CN, CO, and CO₂) were studied. The collision cross sections were calculated from numerous experimental molecular-beam scattering data. The bifurcation approximation to the binary diffusion coefficients and the Wilke-Saxena approximate techniques were effectively used to simplify numerical algorithms of heat transfer calculations.

Nomenclature

- = Sonine expansion coefficients, Eqs. (2) and (3) a_{ip}
- approximation parameter, $1.2A^* \approx 1.452$, Eq. (18) b =
- $b_{ip} c_{ip}^{ji}$ Sonine expansion coefficients, Eq. (4) =
- = Sonine expansion coefficients, Eq. (1)
- heat capacity of species i at constant pressure =
- C_{pi} C₁ approximation coefficient, -0.5, Eq. (22) =
- = heat capacity of species *i* at constant volume
- approximation parameter, Eq. (17) -
- c_{vi} D D_i D_{ik} energy diffusion coefficients, Eq. (11) =
- mass diffusion coefficients, Eq. (1)
- thermal diffusion coefficients, Eq. (2)
- D_i F_i = bifurcation parameter for species i
- f_i $I^{(l,s)}$ = n_i/n , mole fraction of species *i*
- tabulated integrals, Eq. (16) =
- k Boltzmann constant
- М = average molecular weight
- = molecular mass of species i m_i
- N = total number of species
- = average specific number of molecules n
- = pressure p
- q matrix elements of the Chapman-Enskog = approximation, Eqs. (5) and (7)
- R distance between the centers of mass of the = interacting molecules
- R = universal gas constant
- = interatomic distance
- Т translational temperature
- coefficients of binary diffusion ${\mathcal T}_{ij}$ =
- V interaction potential function, Eq. (14)
- approximation parameters, Eq. (22) Z_i
- $= \rho_i / \rho$, mass-concentration of species *i* α_i
- β parameter of the interaction potential, Eq. (14) =
- $egin{array}{c} \gamma \ \delta_{ij} \end{array}$ approximation parameter, Eq. (16)
- Kronecker delta symbols =
- viscosity coefficient η =
- λ = thermal conductivity coefficient
- number of terms in the Sonine polynomial series ξ =
- = density

- σ_{ij} = collision diameter of two molecules i, j
- = correction factor, Eq. (11)
- $\hat{\Omega}_{ii}^{(l,s)*} = \text{collision integrals, Eq. (16)}$
- = approximation parameters, Eq. (18) \varkappa_j

Subscripts

= species

- = internal degrees of freedom in
- 0 = monatomic-gas parameter
- 1 = first approximation, Eqs. (12) and (13)

Introduction

"HE transport properties of gas components at high temperatures are important in heat and mass transfer problems of hypersonic vehicle flights in the Earth and Martian atmospheres. This data could be of significant interest in future NASA missions.^{1,2} Methods of calculating transport coefficients of dissociating air at high temperatures were developed (see Refs. 3-16). Calculations of the transfer coefficients both by kinetic theory¹⁷ and by approximate methods^{12,18-20} are based on data regarding the elastic interaction potentials among mixture components. Experimental investigation of the coefficients has been restricted to the range of temperature $T \le 2000 \text{ K.}^{10,17,21-24}$ At a higher temperature the transport coefficients have only been measured in several unique experiments (i.e., see Refs. 4, 9, 16, and 25-28). Quantum-mechanical calculations of elastic-interaction processes are only possible for molecules with a simple electron structure²⁹ as well as for nitrogen and oxygen atoms in their ground states.^{13,15} The major sources of information on the interaction of molecules at high temperatures are data of molecular-beam scattering experiments.^{6-8,10} Different models of intermolecular potentials (the exponential repulsive potential^{3,7,10,19,20} and the inverse power attractive and repulsive potentials^{6-8,11,30,31}) have been used in these studies. The experimental information on the atom-molecule and molecule-molecule interactions has been also analyzed by Sokolova,³⁰ Jordan et al.,³¹ and Gromov and Eremyan.¹

Unfortunately, the experimental data regarding atom-atom interactions are unavailable and different approximation tech-niques should be developed.^{9,10,13,15} When the atom-atom interactions are described by an exponential repulsion, the average molecule-molecule interaction can be approximated by an additive-potential method, proposed by Amdur et al.³² Using this technique, Yun and Mason³ found the approximate correlations between the atom-molecule and molecule-molecule potentials for air species. This method was effectively used by Riabov²⁰ for calculations of transfer coefficients of

Received March 10, 1995; presented as Paper 95-2058 at the AIAA 30th Thermophysics Conference, San Diego, CA, June 19-22, 1995; revision received Aug. 15, 1995; accepted for publication Nov. 6, 1995. Copyright © 1995 by the American Institute of Aeronautics and Astronautics, Inc. All rights reserved.

Visiting Associate Professor, Department of Mechanical Engineering, 100 Institute Road. Member AIAA.

dissociating air with sublimation products of graphite. In the present study, the additive-potential method³² is developed for calculating interaction potentials and transfer coefficients (viscosity, chemically frozen thermal conductivity, multicomponent mass diffusion, and thermal diffusion) for air and gas components of the Martian atmosphere.

The calculations of the transfer coefficients using accurate expressions of the kinetic theory of gases^{17,33} are carried out with a great consumption of computing time. The simplifications of the calculation technique were developed by Wilke,³⁴ Mason and Saxena,³⁵ Gupta et al.,¹⁶ Kendall et al.,¹⁸ and Riabov.^{11,12,20} Considerably simpler transfer-coefficient expressions could be obtained using bifurcational approximation for the binary-diffusion coefficients.^{11,18,20} In the present study, this approximate technique has been developed to calculate transfer coefficients for species of the Martian atmosphere.

Chapman-Enskog Method and Transfer Coefficients

The Chapman-Enskog iteration technique^{13,25} has been used to solve the Boltzmann equation for the case of a quasineutral mixture of nonexcited monatomic gases. At the hydrodynamic stage the solution of the Boltzmann equation is presented as an asymptotic series according to the small parameter Kn, which is the Knudsen number. The expressions for the diffusion and heat fluxes, and the viscous stress tensor through the gradients of species concentrations, temperature, pressure, and the components of strain tensor are given in Refs. 17, 33, 36, and 37.

The calculation of the transfer coefficients is based on the solution of integral equations using the Sonine polynomial series, i.e., $c_{i0}^{ii}(\xi)$, $a_{i0}(\xi)$, $a_{i1}(\xi)$, $b_{i0}(\xi)$.^{7,33,36} The expressions of the coefficients through the Sonine terms are

$$D_{ij} = (\rho n_i / 2nm_j) (2kT/m_i)^{1/2} c_{i0}^{ji}(\xi)$$
(1)

$$D_i^T = (n_i m_i/2) (2kT/m_i)^{1/2} a_{i0}(\xi)$$
(2)

$$\lambda' = -\frac{5}{4} k \sum_{j=1}^{N} n_j \left(\frac{2kT}{m_j}\right)^{1/2} a_{j1}(\xi)$$
(3)

$$\eta = \frac{1}{2} kT \sum_{j=1}^{N} n_j b_{j0}(\xi)$$
 (4)

The approximate magnitudes of the transfer coefficients are functions of a number of valuable nonzero terms in the polynomial series ξ , which could be calculated from the following system of the linear algebraic equations:

$$\sum_{j=1}^{N} \sum_{p=0}^{M} q_{ij}^{mp} a_{jp} = -\frac{15\sqrt{\pi}}{2} n_i \delta_{m1}$$
(5)

$$\sum_{j=1}^{N} \sum_{p=0}^{M} \tilde{q}_{ij}^{mp} b_{jp} = 5n_i \left(\frac{2\pi m_i}{kT}\right)^{1/2} \delta_{m0}$$
(6)

$$\sum_{i=1}^{N} \sum_{p=0}^{M} q_{ij}^{mp} c_{jp}^{hk} = 3\sqrt{\pi} (\delta_{ik} - \delta_{ih}) \delta_{m0}$$
(7)

where indexes are i, h, $k = 1, \ldots, N$; $m = 0, 1, \ldots, M$, $M = \xi - 1$.

The coefficients q_{ij}^{mp} and q_{ij}^{mp} in Eqs. (5-7) could be calculated using bracket integrals.^{33,36} The expressions for the coefficients and integrals are given in Refs. 17, 33, 36, and 38-41. The form of the thermal conductivity coefficient in the case of monatomic gas mixture is^{17,36,39}

$$\lambda_0 = \lambda' + \lambda'', \qquad \lambda'' = \frac{\rho k}{n} \sum_{i+1}^N \sum_{j=1}^N \frac{E_{ij} D_i^T D_j^T}{n_i m_i m_j}$$
(8)

where E_{ii} is the matrix of elements $D_{ji}m_{i}$.

Calculation methods for the thermal conductivity have been analyzed (see Refs. 35 and 42–49). In calculating the thermal conductivity of polyatomic gases, the contribution of the internal degrees of freedom of the molecules should be taken into account by introducing the Eucken correction⁵⁰ in Eq. (8). In the case of $\xi = 1$, the Eucken correction term has been obtained by Galkin et al.^{44–46} The same technique could be applied in the general case at $\xi > 1$.^{11,37} The expression of the thermal conductivity of polyatomic gas mixture is

$$\lambda = \lambda_0 + \sum_{i=1}^N n_i c_{\nu i} D_i \tag{9}$$

In Eq. (9) the diffusion coefficients D_i are calculated using binary diffusion coefficients \mathcal{T}_{ij} , collision integrals $\Omega_{ij}^{(1,1)*}$, and correction terms $\psi_i^{(M)}$ as follows:

$$\mathcal{T}_{ij} = \frac{3}{16n} \sqrt{\frac{2\pi kT(m_i + m_j)}{m_i m_i}} \frac{1}{\pi \sigma_{ij}^2 \Omega_{ij}^{(1,1)*}}$$
(10)

$$D_{i} = \left(\sum_{j=1}^{N} \frac{f_{j}}{\mathcal{T}_{ij}}\right)^{-1} \frac{1}{[1 - \psi_{i}^{(M)}]}$$
(11)

The correction terms^{11,37} $\psi_i^{(M)}$ are functions of ξ and bracket integrals.³⁶ Nonelastic collisions have been evaluated using the Mason–Monchick approximate method.⁴⁷⁻⁴⁹ The Brokaw effect,⁵¹ which is significant in the analysis of the reactive thermal conductivity for a flow in chemical equilibrium, has not been calculated.

Transfer coefficients could be obtained using the technique described previously by the Gauss method of the solution of the linear algebraic equation system (5–7). The number of arithmetic calculations should be approximately equal to $(\xi \times N)^3$. As a result, a significant amount of calculations per one set of the gasdynamic parameters should be made. To reduce computational time different approximate methods of computing the transfer coefficients have been developed. Wilke³⁴ developed a simple method to estimate the viscosity coefficient of a gas mixture:

$$\eta_{1} = \sum_{i=1}^{N} \eta_{i} \left(1 + \sum_{k=1,k\neq i}^{N} G_{ik} \frac{f_{k}}{f_{i}} \right)^{-1}$$

$$\eta_{ik} = \frac{5}{16} \frac{(2\pi m_{ik} kT)^{1/2}}{\pi \sigma_{ij}^{2} \Omega_{ij}^{(2,2)*}}, \qquad G_{ik}^{\eta} = \frac{2m_{k}}{m_{i} + m_{k}} \frac{\eta_{i}}{\eta_{ik}}$$
(12)

A similar expression for the thermal conductivity coefficient was proposed by Mason and Saxena³⁵:

$$\lambda_{1} = \sum_{i=1}^{N} \lambda_{i}' \left(1 + \sum_{k=1,k\neq i}^{N} G_{ik}^{\lambda} \frac{f_{k}}{f_{i}} \right)^{-1}, \qquad G_{ik}^{\lambda} = L_{ik} \frac{\lambda_{i}}{\lambda_{ik}}$$
$$\lambda_{i}' = \lambda_{0i} \left(0.115 + 0.354 \frac{c_{pi}m_{i}}{k} \right), \qquad \lambda_{0i} = \frac{15}{4} \eta_{i} \frac{k}{m_{i}} \quad (13)$$
$$L_{ik} = \frac{\left[1.25(6m_{i} - 5m_{k}) + 3m_{k}B_{ik}^{*} \right](m_{i} - m_{k}) + 8m_{i}m_{k}A_{ik}^{*}}{2A_{i}^{*}(m_{i} + m_{k})^{2}}$$

Parameters G_{ik}^{λ} and G_{ik}^{η} were analyzed by Ferziger and Kaper³³ in detail. The computational results using Eqs. (12) and (13) correlate well with experimental data^{23,34} for viscosity of binary and triple mixtures under room temperatures and for thermal conductivity under the conditions $T \leq 1000$ K. The applicability of the approximate formulas (12) and (13) for equilibrium dissociating air was considered by Gupta et al.¹⁶ and Riabov.^{11,12} Other approximations for the thermal conductivity coefficient were analyzed by Ferziger and Kaper,³³ Ma-

son and Monchick,⁴⁷ and Lebed and Riabov.⁴⁸ These techniques require a small amount ($\sim N^2$) of calculations.

Interaction Potentials

Numerous experimental data on the molecule-molecule and atom-molecule interactions were obtained (see Refs. 6, 7, 10, 14, 31, and 32). The first investigations^{8,30,32} were based on the assumption of an inverse power dependence of the interaction potential $V = K/R^s$ on the distance between the centers of mass of the molecules R. The analysis of the experimental conditions provided by Leonas⁷ and Cubley and Mason¹⁰ shows that the parameter s is not constant. They recommended the use of repulsive potential in the form:

$$V = V_0 \exp(-\beta R) \tag{14}$$

 Table 1
 Potential parameters for interactions of air species

Interaction	Coefficient V_0 , eV	Coefficient β , Å ⁻¹	Source
0-0	1410	4.14	Cubley and Mason ¹⁰
O-N	348.2	3.41	Cubley and Mason ¹⁰
$O - O_2$	4530	4.039	Riabov ^{20 a}
$O - N_2$	860.4	3.331	Riabov ^{20 a}
O-NO	2142	3.717	Riabov ^{20 a,b}
N-N	86.0	2.68	Cubley and Mason ¹⁰
$N-O_2$	905.7	3.32	Riabov ^{20 b}
$N-N_2$	184.9	2.614	Riabov ^{20 b}
N-NO	428.6	2.983	Riabov ^{20 b}
$O_2 - O_2$	1.485×10^{4}	3.964	Riabov ^{20 a}
$O_2 - N_2$	2316	3.267	Riabov ^{20 b}
$O_2 - NO$	6373	3.644	Riabov ^{20 a,b}
$N_2 - N_2$	415.7	2.573	Riabov ^{20a}
$N_2 - NO$	52.38	1.761	Riabov ²⁰
NO-NO	2678	3.303	Riabov ^{20 a,b}

*Data correlate at high accuracy with data of Cubley and Mason.¹⁰ *Data correlate at high accuracy with data of Leonas.⁷

 Table 2
 Potential parameters for interactions of Martian atmospheric species

T	Coefficient	Coefficient	0
Interaction	<i>v</i> ₀ , ev	р, А	Source
C–C	348	3.37	Gromov and Eremyan ¹⁹
C-0	700.5	3.755	Riabov ²⁰
C-N	173	3.025	Riabov ²⁰
CO2	1997	3.659	Riabov ²⁰
$C-N_2$	398.5	2.956	Riabov ²⁰
C-NO	947.6	3.33	Riabov ²⁰
C-CN	587.4	3.109	Riabov ²⁰
C-CO	1368	3.496	Riabov ²⁰
C-CO ₂	4843	3.5	Riabov ²⁰
N-CN	271.5	2.771	Riabov ²⁰
N-CO	615.7	3.149	Riabov ^{20 a}
N-CO ₂	1855	3.156	Riabov ²⁰
O-CN	1302	3.488	Riabov ²⁰
0-C0	3095	3.882	Riabov ^{20 a}
O-CO ₂	1.31×10^{4}	3.879	Riabov ²⁰
O ₂ -CN	3600	3.416	Riabov ²⁰
O ₂ -CO	9683	3.808	Riabov ²⁰
$O_2 - CO_2$	4.268×10^{4}	3.822	Riabov ^{20 a}
N ₂ -CN	621.7	2.724	Riabov ²⁰
$N_2 - CO$	22.17	1.432	Riabov ²⁰
$N_2 - CO_2$	4738	3.118	Riabov ^{20 a}
NO-CN	1568	3.084	Riabov ²⁰
NO-CO	4075	3.473	Riabov ^{20 a}
NO−CO₂	1.613×10^{4}	3.51	Riabov ²⁰
CN-CN	950.6	2.879	Riabov ²⁰
CN-CO	2371	3.253	Riabov ²⁰
CN-CO ₂	7868	3.268	Riabov ²⁰
CO-CO	6280	3.646	Riabov ²⁰ *
CO-CO ₂	2.494×10^{4}	3.654	Riabov ²⁰ *
$CO_2 - CO_2$	1.049×10^{5}	3.666	Riabov ^{20 a}

^{*}Data correlate at high accuracy with data of Leonas.⁷

This approximation has been successfully used to describe the interaction of inert gases⁷ and of identical atoms.^{9,10} The additive-potential method, proposed by Amdur et al.³² for approximate calculations of V(R), gives acceptable accuracy in calculating the parameters of the atom-molecule and molecule-molecule interactions.^{3,10} According to this method, the interaction potential of the molecules AB and CD is written as^{3,7}

$$V(R) = V(r_{\rm AC}) + V(r_{\rm BC}) + V(r_{\rm AD}) + V(r_{\rm BD})$$
(15)

The potential in Eq. (15), averaged over equiprobable orientations, determines the effective spherically symmetric potential corresponding to the point force center. This approximating procedure was described in detail by Yun and Mason³ and Cubley and Mason.¹⁰ The combinatorial rule,³ $V_{ij} = (V_{ii}V_{jj})^{1/2}$, is usually employed to make up the currently unknown data on the interaction potential.

Using the collision integrals $\Omega_{ij}^{(i,s)*}$ given in Refs. 10, 17, 33, 36, and 52, the transfer coefficients of multicomponent gas mixture may be calculated by the method described previously. According to Monchick,⁵² the expressions for collision integrals calculated for the exponential repulsive potential [see Eq. (14)] take the form:

$$\Omega^{(l,s)*} = \frac{8\gamma^2 I^{(l,s)}(\gamma)}{\beta^2 (s+1)! \left[1 - \frac{1}{2} \frac{1 + (-1)^l}{1 + l}\right] \sigma^2}, \qquad \gamma = \ell n \left(\frac{V_0}{kT}\right)$$
(16)

The functions $I^{(l,s)}(\gamma)$ for various l, s, and values $3.5 \le \gamma \le 28.5$ were tabulated by Monchick.⁵² In the present study, these functions are approximated by cubic splines.

In this study the additive-potential method has been applied to calculate the dependencies V(R) by the previous method for pairs of species O, N, O₂, N₂, NO, C, CN, CO, CO₂, which are the major components of dissociating gases in the Earth and Martian atmospheres.^{1,2} The distance between the centers of mass of the molecules was varied in the range $0.6 \le R \le$ 5.5 Å. The results are approximated by an exponential function, using the least-squares method.²⁰ The basic functions adopted are the atom-atom interaction exponential potentials, the parameters of which are taken from Refs. 10 and 19. Data on the molecular structure are taken from Ref. 53.

The potential parameters V_0 and β obtained in the present study are shown in Table 1 for dissociating air and in Table 2



Fig. 1 Interaction potentials of N atoms with molecules: $N-N_2$ (curves 1-3), $N-O_2$ (curves 4-6), N-NO (curves 7-9). Comparison of the present calculations (solid lines 3, 6, and 9) and experimental data of Leonas⁷ (dashed lines 1, 4, and 7) and Cubley and Mason¹⁰ data (dot-dashed lines 2, 5, and 8).



Fig. 2 Interaction potentials of CO_2 molecules with molecules of the Martian atmospheric gases: CO_2-N_2 (curves 1 and 2), CO_2-O_2 (curves 3 and 4), CO_2-CO (curves 5 and 6), and CO_2-CO_2 (curves 7 and 8). Comparison of the present calculations (solid lines 2, 4, 6, and 8) and experimental data of Leonas⁷ (dashed lines 1, 3, 5, and 7).

for dissociating Martian atmospheric gases. The calculated interaction potentials are compared with the data of Leonas⁷ and Cubley and Mason¹⁰ (see Figs. 1 and 2). The agreement of the results is basically good (see notes in Tables). The additivepotential method used here was extremely important to complete the information on the interaction potentials, especially with the participation of atoms of carbon and oxygen.

Calculation of Transport Coefficients

For dissociating gas components in the Earth and Mars atmospheres at temperatures from 2000 to 10,000 K, the collision integrals $\Omega_{ij}^{(I,s)*}$ are calculated, as well as the viscosity η_i and binary-diffusion coefficients, according to Eqs. (16), (10), and (12). The best agreement of the present results on $\Omega_{ij}^{(I,s)*}$ for air is with the data of Leonas,⁷ Cubley and Mason,¹⁰ and Gromov and Eremyan.¹⁹ The present data for interactions with participation of carbon correlate well with the results of Leonas,⁷ Leonas and Rodionov,¹⁴ Gromov and Eremyan,¹⁹ and Riabov.²⁰ Here the values of the collision integrals with the parameters in Tables 1 and 2 are intermediate between the results of other studies.^{7,19} The difference of the results may reach 25% in the given temperature range (i.e., see Fig. 3). Parameters of the collision-integral ratio^{17,33} are approximately constant, i.e., $A^* \approx 1.21$ and $B^* \approx 1.17$.

The viscosity coefficients η_i for air components and species of the Martian atmosphere are shown in Figs. 4 and 5, correspondingly. These data are in good agreement (about 10%) with the data for O₂, N₂, NO, and CO₂ from Refs. 21–24, 52, and the data¹⁵ for O atoms. The calculated viscosity coefficients of equilibrium dissociating oxygen at constant pressures of 0.1 and 1 atm are shown in Fig. 6. These values are in good agreement with experimental data of Hartunian and Marrone.²⁵

Further analysis of the transfer coefficients is made for equilibrium dissociating air in the range of temperature from 2000 to 10,000 K, and pressure 1 atm, where the concentration of ions and electrons is negligible. Air-component concentrations were calculated by Predvoditelev et al.^{54,55} The results were obtained for exponential repulsion potentials (see Table 1).

The viscosity and chemically frozen thermal conductivity coefficients calculated by the Chapman-Enskog method at $\xi = 1$ [Eqs. (1-11)] are shown in Figs. 7 and 8, correspondingly (filled squares). The results of calculations using the approximate formulas (12) and (13) are presented in Figs. 7 and 8 as empty squares. From comparison of the results, it is found that the approximate formulas provide data with an error of 1% for viscosity and 2% for thermal conductivity. The influ-

ence of the higher order of approximation at $\xi = 2$ on the coefficients is insignificant (less than 1%).

The coefficients of multicomponent mass diffusion of molecular nitrogen with other components at pressure p = 1 atm (solid lines) and 0.1 atm (dashed lines) are shown in Fig. 9. The influence of pressure is significant at $T \ge 3500$ K, where the level of dissociation is high. The values of multicomponent-diffusion coefficients of atoms is larger by approximately



Fig. 3 N_2-N_2 transport collision integrals $\sigma^2 \Omega_{ij}^{(2,2)*}$ (solid lines) and $\sigma^2 \Omega_{ij}^{(1,1)*}$ (dashed lines). Comparison of the present calculations (**T**, **D**) and data of Cubley and Mason¹⁰ (**A**, \triangle) and Gupta et al.¹⁶ data (**0**, \bigcirc).



Fig. 4 Viscosity coefficients η_i for air components vs temperature *T*. Comparison of the present calculations and data of Levin et al.¹⁵ (∇ , O atoms), and Monchick⁵² data (\Box , N₂).



Fig. 5 Viscosity coefficients η_i for components of the Martian atmosphere vs temperature T.



Fig. 6 Viscosity of equilibrium dissociating oxygen at pressures of 0.1 and 1 atm. Comparison of the present calculation results and the experimental data of Hartunian and Marrone²⁵ (\blacksquare), and the data of Yun et al.⁴ (\Box).



Fig. 7 Viscosity coefficient η for equilibrium air vs temperature T at pressure p = 1 atm. The present results (**I**) are compared with the Wilke approximation³⁴ (**I**) and the bifurcation approximation (\blacktriangle).

the factor of 2 in the total range of temperature and pressure. This phenomenon correlates with the difference in the viscosity of the atomic and molecular species (see Fig. 4).

The absolute values of thermal-diffusion coefficients at $\xi = 2$ and pressure of 1 atm are shown in Fig. 10. The coefficients are negative for atoms at $T \le 7000$ K and for molecules of N₂ and NO at $T \le 2500$ K. The dissociation is the major factor in decreasing the thermal-diffusion coefficients of molecules. These coefficients were taken into account to estimate the magnitude of the parameter λ'' in Eq. (8). This parameter is less than 1% of the total value of λ . This fact is in a good agreement with the results of Muckenfuss and Curtiss.⁵⁶

The transport coefficients of the Martian atmospheric gases have been analyzed under the thermodynamic equilibrium conditions in the range of temperature from 2000 to 6000 K, and pressure 1 atm. In accordance with the study of Mitcheltree and Groffo,⁵⁷ we assumed that the Martian undisturbed atmosphere consists of 97% CO₂ and 3% N₂. The equilibrium constants for chemical reactions between nine species were taken from Ref. 57. The thermodynamic functions of species were calculated using data from JANAF tables.⁵⁸

The viscosity and chemically frozen thermal conductivity coefficients calculated by the Chapman-Enskog method at $\xi = 2$ are shown in Figs. 11 and 12, correspondingly (filled squares). The results of calculations using the approximate formulas (12) and (13) are presented in Figs. 11 and 12 as empty squares. From comparison of the results, it is found that the



Fig. 8 Coefficient of chemically frozen thermal conductivity λ' for equilibrium air vs temperature *T* at pressure p = 1 atm. The present results (**■**) are compared with the Mason–Saxena approximation³⁵ (**□**) and the bifurcation approximation (**▲**).



Fig. 9 Multicomponent mass-diffusion coefficients ρD_{ij} for equilibrium air vs temperature T at pressure p = 1 atm (solid lines) and p = 0.1 atm (dashed lines).

approximate formulas provide data with an error of 1% for viscosity and 3% for thermal conductivity. The absolute values of the thermal-diffusion coefficients of species having the biggest values of concentration are shown in Fig. 13.

Approximate Method of Calculations of Transfer Coefficients

As mentioned in the Introduction, approximate methods of estimating transfer coefficients should be developed. Considerably simpler expressions than Eqs. (1-9) could be obtained using the bifurcational approximation for the binary-diffusion coefficients.^{12,18,59,60} According to this technique, the binary-diffusion coefficients could be approximated by the form

$$\mathcal{T}_{ii} = D/F_i F_j \tag{17}$$

where D = D(p, T) is the parameter of the multicomponent mixture and $F_i = F_i(T)$ is the parameter of the species *i*. The possibility of using the correlation relation in Eq. (17) was analyzed by Kendall et al.,¹⁸ Bartlett et al.,⁵⁹ Rolin et al.,⁶⁰ and Riabov^{12,20} for various mixtures of complex composition. The results of the present study also indicate high accuracy of the correlation in Eq. (17) for dissociating gases in the atmospheres of Earth and Mars.

The values of F_i for these mixtures are found by the leastsquares method for the binary-diffusion coefficients calculated



Fig. 10 Thermal diffusion coefficients D_i^T for equilibrium air vs temperature T at pressure p = 1 atm. The present results (solid lines) are compared with the bifurcation approximation (dashed lines).



Fig. 11 Viscosity coefficient η for equilibrium dissociating gases of the Martian atmosphere vs temperature T at pressure p = 1 atm. The present results (**■**) are compared with the Wilke approximation³⁴ (**□**) and the bifurcation approximation (**▲**).

from Eq. (10). The values of D(p, T) for molecular oxygen were taken as the normalizing quantities ($F_{O_2} = 1$). The parameters F_i for the system of elements O-N-C with an exponential repulsive potential are shown in Fig. 14. The magnitudes of F_i depend weakly (with the accuracy of 3%) on the temperature in the range 2000 $\leq T \leq 10,000$ K. The mean relative error is no more than 3% and the maximum error of the complex $\rho T_{ij}/M$ is 11.5%.²⁰ This bifurcational model leads to a mean absolute error of the value of the parameter $\rho T_{ij}/M$, which is more than a factor of 10 less than for the model of equal diffusion coefficients. This confirms the conclusions of Kendall et al.¹⁸ and Riabov.^{12,20} The use of the approximations by Eq. (17) is entirely valid for engineering applications because of the small number (~N) of calculations.

Following Refs. 12 and 33, the simple algebraic formulas are obtained for η and λ' :

$$\eta = \frac{\rho D \varkappa_2}{\varkappa_1 M b}, \qquad \varkappa_1 = M \sum_{j=1}^N \frac{\alpha_j F_j}{M_j}, \qquad \varkappa_2 = \sum_{j=1}^N \frac{M_j f_j}{F_j} \quad (18)$$

 $\lambda' = \lambda'_0 + \lambda_{\rm in}, \qquad b = 1.2A^* \tag{19}$

$$\lambda'_{0} = \frac{15}{4} \frac{\rho D \Re}{M b} \sum_{j=1}^{N} \frac{f_{j}}{F_{j} Q_{j}}, \qquad Q_{j} = \sum_{k=1}^{N} L_{jk} f_{k} F_{k} \qquad (20)$$



Fig. 12 Coefficient of chemically frozen thermal conductivity λ' for equilibrium dissociating gases of the Martian atmosphere vs temperature *T* at pressure p = 1 atm. The present results (**I**) are compared with the Mason–Saxena approximation³⁵ (**D**) and the bifurcation approximation (\blacktriangle).



Fig. 13 Thermal diffusion coefficients D_i^T for equilibrium dissociating gases of the Martian atmosphere vs temperature T at pressure p = 1 atm. The present results (solid lines) are compared with the bifurcation approximation (dashed lines).

$$\Lambda_{\rm in} = \frac{\rho D \mathcal{R}}{\varkappa_1 M} \sum_{j=1}^N \frac{f_j}{F_j} c_{{\rm in},j}$$
(21)

The thermal-diffusion coefficient is approximated by the expression^{12,59}:

$$D_i^T = c_i \frac{\rho D \varkappa_2}{\varkappa_1 M} (z_i - \alpha_i), \qquad z_i = \frac{M_i f_i}{\varkappa_2 F_i}$$
(22)

The results of calculating the viscosity, frozen thermal conductivity, and thermal-diffusion coefficients of equilibrium-dissociated air and Martian atmospheric gases under considered conditions using bifurcation approximation are presented in Figs. 7, 8, 11, and 12 (dot-dashed lines), and Figs. 10 and 13 (dashed lines). According to the estimates, the difference in the results of accurate and approximate approaches over the whole temperature range is no more than 3% for the viscosity and 8.5% for the chemically frozen thermal conductivity. The approximation by Eq. (22) is helpful for the qualitative analysis of the thermal-diffusion coefficients. This approximation algorithm was successfully used by Riabov and Provotorov,^{61,62} and Riabov and Botin⁶³ in the structure analysis of the thin viscous shock layers near hypersonic vehicles and near probes in supersonic hydrogen-combustion zones.



Fig. 14 Parameters F_i vs temperature T.

Concluding Remarks

The obtained results indicate the possibility of applying the approximate bifurcation method and the Wilke-Mason-Saxena technique³³⁻³⁵ to calculate the transfer coefficients of mixtures of complex composition. The accuracy of these techniques is acceptable for practical applications. This method could be effectively used for solving heat transfer problems in future planetary missions.

Acknowledgments

The author would like to express gratitude to V. S. Galkin, M. N. Kogan, and I. V. Lebed for their fruitful participation in developing methods for solving the problem.

References

¹Park, C., "Review of Chemical-Kinetic Problems of Future NASA Missions, I: Earth Entries," *Journal of Thermophysics and Heat Transfer*, Vol. 7, No. 3, 1993, pp. 385–398.

²Park, C., Howe, J. T., Jaffe, R. L., and Candler, G. V., "Review of Chemical-Kinetic Problems of Future NASA Missions, II: Mars Entries," *Journal of Thermophysics and Heat Transfer*, Vol. 8, No. 1, 1994, pp. 9–23.

³Yun, K. S., and Mason, E. A., "Collision Integrals for the Transport Properties of Dissociating Air at High Temperatures," *Physics of Fluids*, Vol. 5, No. 4, 1962, pp. 380–386.

⁴Yun, K. S., Weissman, S., and Mason, E. A., "High-Temperature Transport Properties of Dissociating Nitrogen and Dissociating Oxygen," *Physics of Fluids*, Vol. 5, No. 6, 1962, pp. 672–678.

⁵Yos, J. M., 'Transport Properties of Nitrogen, Hydrogen, Oxygen and Air to 30,000 K,' AVCO Corp., TM RAD-TM-63-7, March 1963. 'Belyaev, Y. N., Kamyshov, N. V., Leonas, V. B., and Sermyagin,

A. V., "The Experimental Investigation of Short-Range Repulsive Forces Between Atoms and Molecules of Atmospheric Gases," *Entropie*, Vol. 30, Nov.–Dec. 1969, pp. 173–177.

⁷Leonas, V. B., "Studies of Short-Range Intermolecular Forces," Soviet Physics Uspekhi, Vol. 15, No. 3, 1973, pp. 266–281.

⁸Sokolova, I. A., "Transport Coefficients of Air in the Temperature Range from 3,000 K to 25,000 K and Pressures 0.1, 1, and 100 Atm," *Journal of Applied Mechanics and Technical Physics*, Vol. 14, No. 2, 1973, pp. 20–32.

⁹Capitelli, M., and Devoto, R. S., "Transport Coefficients of High-Temperature Nitrogen," *Physics of Fluids*, Vol. 16, No. 11, 1973, pp. 1835-1841.

¹⁰Cubley, S. J., and Mason, E. A., "Atom-Molecule and Molecule-Molecule Potentials and Transport Collision Integrals for High-Temperature Air Species," *Physics of Fluids*, Vol. 18, No. 9, 1975, pp. 1109-1111.

¹¹Riabov, V. V., "Calculation of Transport Coefficients of Equilibrium Dissociated Air," *Trudy TsAGI*, No. 2045, 1980, pp. 3-14 (in Russian).

Russian). ¹²Riabov, V. V., "Approximate Calculation of Transport Coefficients in Multicomponent Mixtures," *Journal of Engineering Physics*, Vol. 44, No. 2, 1983, pp. 183–189.

¹³Rainwater, J. C., Biolsi, L., Biolsi, K., and Holland, P. M., "Transport Properties of Ground State Nitrogen Atoms," *Journal of Chemical Physics*, Vol. 73, No. 3, 1983, pp. 1462–1468. ¹⁴Leonas, V. B., and Rodionov, I. D., "High-Energy Scattering of Atoms and Molecules," *Soviet Physics Uspekhi*, Vol. 28, No. 5, 1985, pp. 332-348.

pp. 332-348. ¹⁵Levin, E., Partridge, H., and Stallcop, J. R., "Collision Integrals and High-Temperature Transport Properties for N-N, O-O, and N-O," *Journal of Thermophysics and Heat Transfer*, Vol. 4, No. 4, 1990, pp. 469-477.

¹⁶Gupta, R. N., Yos, J. M., Thompson, R. A., and Lee, K.-P., "A Review of Reaction Rates and Thermodynamic and Transport Properties for an 11-Species Air Model for Chemical and Thermal Nonequilibrium Calculations to 30,000 K," NASA Reference Publication 1232, Aug. 1990.

¹⁷Hirschfelder, J. O., Curtiss, C. F., and Bird, R. D., *Molecular Theory of Gases and Liquids*, Wiley, New York, 1964.

¹⁸Kendall, R. M., Rindal, R. A., and Bartlett, E. P., "A Multicomponent Boundary Layer Chemically Coupled to an Ablating Surface," *AIAA Journal*, Vol. 5, No. 6, 1967, pp. 1063–1071.

¹⁹Gromov, V. G., and Eremyan, N. A., "Investigation of Transport Processes in the High-Temperature Boundary Layer on an Ablating Graphite Surface," *Fluid Dynamics*, Vol. 18, No. 3, 1983, pp. 416– 421.

²⁰Riabov, V. V., "Transfer Coefficients of Multicomponent Air with Sublimation Products of Graphite," *Journal of Engineering Physics*, Vol. 55, No. 1, 1988, pp. 786–791.

²¹Ellis, C. P., and Raw, C. J. G., "High-Temperature Gas Viscosities. II Nitrogen, Nitric Oxide, Boron Trifluoride, Silicon Tetrafluoride, and Sulfur Hexafluoride," *Journal of Chemical Physics*, Vol. 30, No. 2, 1959, pp. 574–576.
 ²²Weissman, S., "Estimation of Diffusion Coefficients from Vis-

²²Weissman, S., "Estimation of Diffusion Coefficients from Viscosity Measurements: Polar and Polyatomic Gases," *Journal of Chemical Physics*, Vol. 40, No. 11, 1964, pp. 3397–3406.

²³Vargaftik, N. B., Handbook on the Thermophysical Properties of Gases and Liquids, Nauka, Moscow, 1972 (in Russian).

²⁴Lavushchev, A. V., and Lyusternik, V. E., "Experimental Determination of Viscosity of Nitrogen up to 2000 K," *High Temperature*, Vol. 16, No. 1, 1978, pp. 172, 173.

²⁵Hartunian, R. A., and Marrone, P. V., "Viscosity of Dissociated Gases from Shock-Tube Heat-Transfer Measurements," *Physics of Fluids*, Vol. 4, No. 5, 1961, pp. 535–543.

²⁶Morris, J. C., Rudis, R. P., and Yos, J. M., "Measurements of Electrical and Thermal Conductivity of Hydrogen, Nitrogen, and Argon at High Temperatures," *Physics of Fluids*, Vol. 13, No. 3, 1970, pp. 608-617.

²⁷Schreiber, P. W., Hunter, A. M., and Benedetto, K. R., "Argon and Nitrogen Plasma Viscosity Measurements," *Physics of Fluids*, Vol. 14, No. 12, 1971, pp. 2696–2702.

²⁸Schreiber, P. W., Hunter, A. M., and Benedetto, K. R., "Measurement of Nitrogen Plasma Transport Properties," *AIAA Journal*, Vol. 10, No. 5, 1972, pp. 670–674.

²⁹McWeeny, R., *Methods of Molecular Quantum Mechanics*, 2nd ed., Academic, London, 1989. ³⁰Sokolova L.A., ⁴¹Tetamola, G. ⁴¹Tetamola,

³⁰Sokolova, I. A., 'Integrals for Collision Components of Partially Ionized Air,'' *Proceedings of the 3rd All-Union Conference on the Physics of Low-Temperature Plasmas*, Moscow State Univ., Moscow, 1971, pp. 121–123 (in Russian).

³¹Jordan, J. E., Colgate, S. O., Amdur, I., and Mason, E. A., "Scattering of High-Velocity Neutral Particles. XVII. Ar-O₂, Ar-N₂, Ar-CO," *Journal of Chemical Physics*, Vol. 52, No. 3, 1970, pp. 1143– 1149.

³²Amdur, I., Longmire, M. S., and Mason, E. A., "Scattering of High-Velocity Neutral Particles. XII. He-CH₄, He-CF₄, CH₄-CH₄ and CF₄-CF₄ Interactions," *Journal of Chemical Physics*, Vol. 35, No. 3, 1961, pp. 895–898.

 ³³Ferziger, J., and Kaper, H. G., Mathematical Theory of Transport Processes in Gases, North-Holland, Amsterdam, 1972.
 ³⁴Wilke, C. R., "A Viscosity Equation for Gas Mixtures," Journal

³⁴Wilke, C. R., "A Viscosity Equation for Gas Mixtures," *Journal of Chemical Physics*, Vol. 18, No. 4, 1950, pp. 517–519.
 ³⁵Mason, E. A., and Saxena, S. C., "Approximation Formula for the second second

³⁵Mason, E. A., and Saxena, S. C., 'Approximation Formula for the Thermal Conductivity of Gas Mixtures,' *Physics of Fluids*, Vol. 1, No. 5, 1958, pp. 361–369.

³⁶Chapman, S., and Cowling, T. G., *The Mathematical Theory of Non-Uniform Gases*, 3rd ed., Cambridge Univ. Press, London, 1970.
 ³⁷Riabov, V. V., "Approximate Calculation of Transport Coeffi-

³⁷Riabov, V. V., "Approximate Calculation of Transport Coefficients of the Earth and Mars Atmospheric Dissociating Gases," AIAA Paper 95-2058, June 1995.

³⁸Tirsky, G. A., "Equations of Motion of Partially Ionized Multicomponent Gas Mixtures in Normal Cauchy Form with Exact Transport Coefficients," *Trudy Instituta Mekhaniki, Moscow State University*, edited by S. S. Grigoryan, No. 32, 1974, pp. 6–22 (in Russian). ³⁹Devoto, R. S., "Transport Properties of Ionized Monatomic Gases," *Physics of Fluids*, Vol. 9, No. 6, 1966, pp. 1230-1240.

⁴⁰Li, C. P., and Devoto, R. S., "Fifth and Sixth Approximations to the Electron Transport Coefficients," *Physics of Fluids*, Vol. 11, No. 2, 1968, pp. 448-450.

⁴¹Kolesnikov, A. F., and Tirsky, G. A., "Hydrodynamics Equations of Partially Ionized Multicomponent Gas Mixtures with Higher Approximation Transport Coefficients Consideration," *Molecular Gasdynamics*, edited by V. V. Struminsky, Nauka, Moscow, pp. 20–44 (in Russian).

⁴²Mason, E. A., Vanderslice, J. T., and Yos, J. M., "Transport Properties of High-Temperature Multicomponent Gas Mixtures," *Physics* of *Fluids*, Vol. 2, No. 6, 1959, pp. 688–694.

⁴³Butler, J. N., and Brokaw, R. S., "Thermal Conductivity of Gas Mixtures in Chemical Equilibrium," *Journal of Chemical Physics*, Vol. 26, No. 6, 1957, pp. 1636–1643.

⁴⁴Galkin, V. S., and Kogan, M. N., "On the Equations for Nonequilibrium Polyatomic Gas Flows in the Eucken Approximation," *The Problems of Hydrodynamics and Continuum Mechanics*, Nauka, Moscow, 1969, pp. 37-52 (in Russian).

⁴⁵Galkin, V. S., Kogan, M. N., and Makashev, N. K., "The Chapman-Enskog Generalized Method," *Doklady Akademii Nauk SSSR*, Vol. 220, No. 2, 1975, pp. 304–307 (in Russian).

⁴⁶Galkin, V. S., and Makashev, N. K., "Kinetic Derivation of the Gas-Dynamic Equation for Multicomponent Mixtures of Light and Heavy Particles," *Fluid Dynamics*, Vol. 29, No. 1, 1994, pp. 140–155.

⁴⁷Mason, E. A., and Monchick, L., "Heat Conductivity of Polyatomic and Polar Gases," *Journal of Chemical Physics*, Vol. 36, No. 6, 1962, pp. 1622–1639.

⁴⁸Lebed, I. V., and Riabov, V. V., "Rotational Relaxation Time and Transfer Coefficients in a Diatomic Gas," *Journal of Applied Mechanics and Technical Physics*, Vol. 24, No. 4, 1983, pp. 447– 454.

⁴⁹Riabov, V. V., ''Nonequilibrium Gas Dynamic Equations and Transfer Coefficients in Diatomic Gas,'' AIAA Paper 94-2400, June 1994.

⁵⁰Eucken, A., "Uber der Wärmeleitvermogen, die spezifische Wärme und die innere Reibung der Gase," Zeitschrift für Physik, Vol. 14, 1913, pp. 324–341.

⁵¹Brokaw, R. S., "Approximate Formulas for the Viscosity and Thermal Conductivity of Gas Mixtures," *Journal of Chemical Physics*, Vol. 29, No. 2, 1958, pp. 391–397.

⁵²Monchick, L., "Collision Integrals for the Exponential Repulsive Potential," *Physics of Fluids*, Vol. 2, No. 6, 1959, pp. 695-700.

⁵³Glushko, V. P., *Thermodynamic Constants of Matter; Handbook in Ten Parts*, Academy of Sciences of the USSR, Moscow, 1982.

⁵⁴Predvoditelev, A. S., Stupochenko, E. V., Samuilov, E. V., Stakhanov, I. P., Pleshanov, A. S., and Rozhdestvenskii, I. B., *Tables of Thermodynamic Functions of Air for the Temperature Range 6,000–* 12,000 K and Pressure Range 0.001–1000 Atm, Infosearch, Ltd., London, 1958.

⁵⁵Predvoditelev, A. S., Stupochenko, E. V., Pleshanov, A. S., Samuilov, E. V., and Rozhdestvenskii, I. B., *Tables of Thermodynamic Functions of Air for the Temperature Range 200–6,000 K and Pressure Range 0.00001–100 Atm*, Computer Center of the U.S.S.R. Academy of Sciences, Moscow, 1962 (in Russian).

⁵⁶Muckenfuss, C., and Curtiss, C. F., "Thermal Conductivity of Multicomponent Gas Mixtures," *Journal of Chemical Physics*, Vol. 29, No. 6, 1958, pp. 1273–1277.

⁵⁷Mitcheltree, R. A., and Gnoffo, P. A., "Wake Flow About a MESUR Mars Entry Vehicle," AIAA Paper 1958, June 1994.

⁵⁸JANAF Thermochemical Tables, 2nd ed., Dow Chemical Co., Midland, MI, July 1970.

⁵⁹Bartlett, E. P., Kendall, R. M., and Rindal, R. A., "A Unified Approximation for Mixture Transport Properties for Multicomponent Boundary-Layer Applications," NASA CR 1063, June 1968.

⁶⁰Rolin, M. N., Yurevich, F. B., and Kondrashov, V. V., "An Approximate Method for Calculating Transfer Coefficients for Partially Ionized Gas Mixtures," *Journal of Engineering Physics*, Vol. 34, No. 3, 1978, pp. 296–302.

⁶¹Riabov, V. V., and Provotorov, V. P., "The Structure of Multicomponent Nonequilibrium Viscous Shock Layers," AIAA Paper 94-2054, June 1994.

⁶²Riabov, V. V., and Provotorov, V. P., "Modeling of Heat Transfer Processes at Catalytic Materials in Shock Tube," *Journal of Thermophysics and Heat Transfer*, Vol. 9, No. 2, 1995, pp. 363–365.

⁶³Riabov, V. V., and Botin, A. V., "Hypersonic Hydrogen Combustion in the Thin Viscous Shock Layer," *Journal of Thermophysics* and Heat Transfer, Vol. 9, No. 2, 1995, pp. 233–239.